IUB-TH-043

Spherically symmetric solutions of a $(4+n)$- dimensional Einstein-Yang-Mills model with cosmological constant

Yves Brihaye*

Faculté des Sciences, Université de Mons-Hainaut, 7000 Mons, Belgium

Betti Hartmann†

School of Engineering and Sciences,
International University Bremen (IUB), 28725 Bremen, Germany

(Dated: February 1, 2008)

arXiv:hep-th/0403296v3 19 Oct 2004

Abstract

We construct solutions of an Einstein-Yang-Mills system including a cosmological constant in 4+n space-time dimensions, where the n-dimensional manifold associated with the extra dimensions is taken to be Ricci flat. Assuming the matter and metric fields to be independent of the n extra coordinates, a spherical symmetric Ansatz for the fields leads to a set of coupled ordinary differential equations. We find that for $n > 1$ only solutions with either one non-zero Higgs field or with all Higgs fields constant and zero gauge field function (corresponding to a Wu-Yang-type ansatz) exist. We give the analytic solutions available in this model. These are "embedded" abelian solutions with a diverging size of the manifold associated with the extra n dimensions. Depending on the choice of parameters, these latter solutions either represent naked singularities or they possess a single horizon.

We also present solutions of the effective 4-dimensional Einstein-Yang-Mills-Higgs-dilaton model, where the higher dimensional cosmological constant induces a Liouville-type potential. The solutions are non-abelian solutions with diverging Higgs fields, which exist only up to a maximal value of the cosmological constant.

PACS numbers: 04.20.Jb, 04.40.Nr, 04.50.+h, 11.10.Kk

---

* Yves.Brihaye@umh.ac.be
† b.hartmann@iu-bremen.de

I. INTRODUCTION

In an attempt to unify electrodynamics and general relativity, Kaluza introduced an extra, a fifth dimension [1] and assumed all fields to be independent of the extra dimension. Klein [2] followed this idea, however, he assumed the fifth dimension to be compactified on a circle of Planck length. The resulting theory describes 4-dimensional Einstein gravity plus Maxwell's equations. One of the new fields appearing in this model is the dilaton, a scalar companion of the metric tensor. In an analogue way, this field arises in the low energy effective action of superstring theories and is associated with the classical scale invariance of these models [3].

Both string theories [4] as well as so-called "brane worlds" [5] assume that space-time possesses more than four dimensions. In string theories, these extra dimensions are -following the idea of Klein- compactified on a scale of the Planck length, while in brane worlds, which assume the Standard model fields to be confined on a 3-brane, they are large or even infinite. It should thus be interesting to study classical solutions of non-abelian gauge theories in higher dimensions and with view to the AdS/CFT [6], respectively dS/CFT [7] correspondence especially including a cosmological constant.

Classical soliton-like solutions of Einstein-Yang-Mills theories in d dimensions where studied in [8]. These solutions are spherically symmetric in the full d dimensions. Solutions with spherical symmetry in only $3+1$ dimensions were considered for $n=1$ codimension in [9], while they were studied for n codimensions in [10]. The manifold associated with the extra dimensions was assumed to be Ricci flat. If all fields are assumed to be independent of the extra coordinates, the $4+n$ dimensional system reduces to an effective 4-dimensional Einstein-Yang-Mills-Higgs-dilaton (EYMHD) system with n independent, massless dilatons and n Higgs triplets. Constraints appear in this model such that only solutions with either one non-zero Higgs field or with all Higgs fields constant and zero gauge field function (Wu-Yang-Type ansatz) exist.

If a cosmological constant is introduced into this model, the effective 4-dimensional action describes an EYMHD system, where now the dilatons become massive due to a Liouville-type potential. The $(4+1)$-dimensional version of this model was studied in [11]. It was found that no asymptotically flat nor de Sitter nor Anti-de Sitter solutions exist in this model- analogue to the Einstein-Maxwell-dilaton case with Liouville potential [12]. Consequently, black hole solutions with unusual asymptotics were constructed, which manifest the "embedded" abelian counterparts of the 4-dimensional solutions found in [13].

In this paper, we study the system of [11] in $4+n$ dimensions, assuming all fields to be independent of the n extra coordinates. In section II, we present the model including the Ansatz, the equations and the analytic solutions, which fufill the constraints that arise in this model. In Section III, we give the 4-dimensional effective action resulting from the model and give analytic solutions, which represent non-abelian solutions with diverging Higgs fields. These solutions exist only up to a maximal value of the cosmological constant. We give our conclusions in Section IV.

II. THE $(4+n)$-DIMENSIONAL Λ-EINSTEIN-YANG-MILLS MODEL

The Λ-Einstein-Yang-Mills Lagrangian in $d=4+n$ dimensions is given by:

$$S=\int\left(\frac{1}{16\pi G_{(4+n)}}\left(R-2\Lambda_{(n+4)}\right)-\frac{1}{4e^2}F^a_{MN}F^{aMN}\right)\sqrt{g^{(4+n)}}d^{(4+n)}x \tag{1}$$

with the SU(2) Yang-Mills field strengths $F^a_{MN}=\partial_M \mathsf{A}^a_N-\partial_N \mathsf{A}^a_M+\epsilon_{abc}\mathsf{A}^b_M \mathsf{A}^c_N$, the gauge index $a=1,2,3$ and the space-time index $M=0,...,(4+n)-1$. $G_{(4+n)}$ and e denote respectively the $(4+n)$-dimensional Newton's constant and the coupling constant of the gauge field theory. $G_{(4+n)}$ is related to the Planck mass M_{pl} by $G_{(4+n)}=M_{pl}^{-(2+n)}$ and e^2 has the dimension of $[\text{length}]^n$. $\Lambda_{(n+4)}$ is the $(4+n)$-dimensional cosmological constant.

In the following, we denote the coordinates $x_{(3+k)}$ by y_k with $k=1,...,n$.

If both the matter functions and the metric functions are independent on y_k, the fields can be parametrized as follows:

$$g^{(4+n)}_{MN}dx^M dx^N=e^{-\Xi}g^{(4)}_{\mu\nu}dx^\mu dx^\nu+\sum_{k=1}^{n}e^{2\zeta_k}(dy^k)^2\ ,\ \ \mu,\nu=0,1,2,3 \tag{2}$$

with

$$\Xi=\sum_{k=1}^{n}\zeta_k \tag{3}$$

and

$$\mathsf{A}^a_M dx^M=\mathsf{A}^a_\mu dx^\mu+\sum_{k=1}^{n}\Phi^a_k dy^k\ . \tag{4}$$

$g^{(4)}$ is the 4-dimensional metric tensor and the ζ_j and Φ^a_j, $j=1,...,n$, play the role of dilatons and Higgs fields, respectively. Note that the factor $e^{2\zeta_k}$ leads to a space-time dependent seize of the manifold associated with the extra dimensions. The same holds true for the factor $e^{-\Xi}$ with respect to the 4-dimensional "physical" manifold.

The case $n = 1$, $\Lambda_5 = 0$ was studied in [9], while the case $n = 1$, $\Lambda_5 \neq 0$ was investigated in [11]. This present paper is an extension of the results in [10], which deals with the case $\Lambda_{n+4} = 0$ for generic n.

A. Spherically symmetric Ansatz

For the metric the spherically symmetric Ansatz in Schwarzschild-like coordinates reads [14]:

$$g^{(4)}_{\mu\nu} dx^\mu dx^\nu = -A^2(r)N(r)dt^2 + N^{-1}(r)dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta d^2\varphi \ , \tag{5}$$

with

$$N(r) = 1 - \frac{2m(r)}{r} \ . \tag{6}$$

For the gauge and Higgs fields, we use the purely magnetic hedgehog ansatz [15] :

$$\mathsf{A}_r{}^a = \mathsf{A}_t{}^a = 0 \ , \tag{7}$$

$$\mathsf{A}_\theta{}^a = (1 - K(r)) e_\varphi{}^a \ , \qquad \mathsf{A}_\varphi{}^a = -(1 - K(r)) \sin\theta e_\theta{}^a \ , \tag{8}$$

$$\Phi^a_j = c_j v H_j(r) e_r{}^a \quad , \quad j = 1, ..., n \ , \tag{9}$$

where v is a mass scale, while c_j are dimensionless constants determining the vacuum expectation values of the Higgs fields $\langle \Phi_j \rangle = c_j v$. In absence of a Higgs potential these have to be set by hand. Finally, the dilatons are scalar fields depending only on r :

$$\zeta_j = \zeta_j(r) \quad , \quad j = 1, ..., n \ . \tag{10}$$

B. Equations of motion

The non-vanishing components of the energy-momentum tensor are given by:

$$\begin{aligned}
T^0_0 &= -e^{\Xi} \left(\sum_{k=1}^{n} (\mathcal{A}_k + C_k) + B + D \right) \ , \\
T^1_1 &= -e^{\Xi} \left(\sum_{k=1}^{n} (-\mathcal{A}_k + C_k) - B + D \right) \ , \\
T^2_2 &= T^3_3 = -e^{\Xi} \left(\sum_{k=1}^{n} \mathcal{A}_k - D \right) \ ,
\end{aligned}$$

$$\begin{aligned} T_j^j &= -e^{\Xi}\left(-2(\mathcal{A}_j + C_j) + \sum_{k=1}^{n}(\mathcal{A}_k + C_k) + B + D\right) \ , \ j = 1, ..., n \ , \\ T_j^k &= e^{\Xi} e^{-(\xi_1+\xi_2)}\left(\frac{N}{2}H_j' H_k' + \frac{K^2}{x^2} H_j H_k\right) \ , \ j \neq k \ , \end{aligned} \tag{11}$$

where we use the abbreviations

$$\mathcal{A}_j = \frac{1}{2} e^{-2\zeta_j} N (H_j')^2 \quad , \quad C_j = e^{-2\zeta_j} K^2 H_j^2 \frac{1}{x^2} \tag{12}$$

and

$$B = e^{\Xi} \frac{1}{x^2} N (K')^2 \quad , \quad D = e^{\Xi} \frac{1}{2x^4} (K^2 - 1)^2 \ . \tag{13}$$

Defining the coupling constant $\alpha = v\sqrt{G_{(4+n)}}$, the rescaled cosmological constant $\Lambda = \frac{1}{e^2 v^2}\Lambda_{(4+n)}$, the mass function $\mu = evm$ and the radial coordinate $x = evr$, we obtain the following differential equations (the prime denotes the derivative with respect to x) :

$$\mu' = \alpha^2 x^2 \left(\sum_{k=1}^{n}(\mathcal{A}_k + C_k) + B + D\right) + \frac{1}{2} N x^2 \Theta + \frac{\Lambda x^2}{4} e^{-\Xi} \tag{14}$$

$$A' = \alpha^2 A x \left(\sum_{k=1}^{n} e^{-2\zeta_k} (H_k')^2 + 2 e^{\Xi} \frac{K'^2}{x^2}\right) + x A \Theta \tag{15}$$

with

$$\Theta \equiv \frac{3}{4}\left(\sum_{k=1}^{n} (\zeta_k')^2 + \frac{2}{3} \sum_{k>k'} \zeta_k' \zeta_{k'}'\right) \ , \tag{16}$$

$$(x^2 A N \zeta_j')' = \alpha^2 A x^2 e^{-\Xi} \left[\frac{2}{n+2} \sum_{M=0}^{d-1} T_M^M - 2 T_j^j\right] - \frac{A x^2}{n+2} \Lambda e^{-\Xi} \quad , \quad j = 1, ..., n \ , \tag{17}$$

$$(e^{\Xi} A N K')' = A\left(\frac{1}{x^2} e^{\Xi} K (K^2 - 1) + \sum_{k=1}^{n} e^{-2\zeta_k} K H_k^2\right) \ , \tag{18}$$

$$(e^{-2\zeta_j} x^2 A N H_j')' = 2 A e^{-2\zeta_j} K^2 H_j \quad , \quad j = 1, ..., n \ . \tag{19}$$

Finally, since the off-diagonal components of the Einstein tensor vanish, we obtain constraints on the fields from the jk-components of the energy-momentum tensor:

$$e^{\Xi} e^{-(\xi_1+\xi_2)} \left(\frac{N}{2} H_j' H_k' + \frac{K^2}{x^2} H_j H_k\right) = 0 \ , \ j \neq k \ . \tag{20}$$

These constraints lead to the conclusion that solutions with either only one non-zero Higgs field or with all Higgs fields constant and zero gauge field function K exist. The former case would effectively correspond to the $4+1$-dimensional model with one Higgs field. In the following, we will discuss the solutions available in $4+n$ dimensions with all Higgs fields being constant.

C. "Embedded" abelian solutions

We assume here that $H_1(x) = H_2(x) = ... \equiv H(x)$ and thus $\zeta_1(x) = \zeta_2(x) = ... \equiv \zeta(x)$.

Note that no solutions with constant seize of the manifold associated with the extra dimensions (i.e. constant dilaton fields) exist in this model, if one chooses either $K(x) = 0$, $H_k(x) = 1$ or $K(x) = 1$, $H_k(x) = 0$. However, solutions with non-constant dilatons exist. Choosing

$$K(x) = 0 \quad , \quad H(x) = 1 \quad , \tag{21}$$

the solutions correspond to "embedded" abelian solutions with magnetic charge $P = 1$ since $F^{(r)}_{\theta\varphi} = \sin\theta$.

The counterparts of these solutions for a model involving one dilaton field with Liouville type potential in 4-dimensional Einstein-Maxwell theory were discussed in [13]. Note that the effective 4-dimensional action which results from our model in the case $n = 1$ [11] corresponds to the 4-dimensional model studied in [13]. Here, however, the effective action contains n dilaton fiels. As was proven in [12], no asymptotically flat nor de Sitter nor Anti-de Sitter solutions exist for the model in [13]. This is also true for the "embedded" abelian counterparts [11] and the system studied here.

Here we present the $(4+n)$-dimensional counterparts of the solutions found in [11]. The solutions read:

$$A(x) = a_0 x^{(n+2)/2} \quad , \quad N(x) = n_0 - n_1 x^{-2(n+1)/n} \quad , \quad \zeta(x) = \zeta_0 + \frac{2}{n}\ln(x) \tag{22}$$

where a_0, n_1 and ζ_0 are not further determined constants, while

$$n_0 = \frac{n^2\alpha^2}{2(n+2)(n+1)} e^{n\zeta_0} - \frac{\Lambda n^2}{4(n+1)(n+2)} e^{-n\zeta_0} \tag{23}$$

and the cosmological constant is given by:

$$\Lambda = (n+2)e^{n\zeta_0} - 2(n+1)\alpha^2 e^{2n\zeta_0} \ . \tag{24}$$

The metric then reads:

$$\begin{aligned} ds^2 &= -e^{-n\zeta_0} a_0^2 x^n \left(n_0 - n_1 x^{-(2n+2)/n}\right) dt^2 + e^{-n\zeta_0} x^{-2} \left(n_0 - n_1 x^{-(2n+2)/n}\right)^{-1} dx^2 \\ &+ e^{-n\zeta_0}\left(d\theta^2 + \sin^2\theta d\varphi^2\right) + e^{2\zeta_0} x^{4/n}\left((dy_1)^2 + ... + (dy_n)^2\right) \end{aligned} \tag{25}$$

The Kretschmann scalar $K = R^{MNOP}R_{MNOP}$ of the solutions is in general very compicated, that's why we give its form here for $n = 2$. It reads:

$$K_{(n=2)} = \frac{2e^{4\zeta_0}}{3}\left(\alpha^4(9e^{12\zeta_0} + 6e^{8\zeta_0} + e^{4\zeta_0}) - \alpha^2(12e^{10\zeta_0} - 4e^{8\zeta_0}) + 4e^{8\zeta_0} + 6 + 18\frac{n_1^2}{x^6}\right) \quad (26)$$

Clearly, the solutions possess a physical singularity at $x = 0$. This is also true for $n \neq 2$, e.g. for $n = 1$ and $n = 3$ the leading divergence is proportional to x^{-10}, respectively $x^{-\frac{10}{3}}$. This clearly indicates that the singularity gets weaker when increasing the number of extra dimensions. We also note that for $n \neq 2$, the Ricci scalar R and the invariant $R^{AB}R_{AB}$ are singular at $x = 0$, e.g. for $n = 1$ the leading divergence is of power x^{-5} and x^{-10} for R, respectively $R^{AB}R_{AB}$. However, for $n = 2$ R and $R^{AB}R_{AB}$ are constant:

$$R_{(n=2)} = 2e^{2\zeta_0}\left(3e^{6\zeta_0}\alpha^2 - 2e^{4\zeta_0} + e^{2\zeta_0}\alpha^2 - 1\right) \ , \quad (27)$$

$$(R^{AB}R_{AB})_{(n=2)} = e^{4\zeta_0}\left(9e^{12\zeta_0}\alpha^4 - 12e^{10\zeta_0}\alpha^2 + 6e^{8\zeta_0}\alpha^4 + 4e^{8\zeta_0} - 4e^{6\zeta_0}\alpha^2 + e^{4\zeta_0}\alpha^4 + 2\right) \quad (28)$$

This can be related to the particular ansatz that we have introduced for the metric (see (2)). Clearly for $n = 2$, the prefactor in front of the 4-dimensional, spherically symmetric part becomes $e^{-2\zeta}$, while the prefactor in front of the extra dimensional part is always $e^{2\zeta}$. For $n = 2$ we thus have $g_{tt} \propto g_{xx}$.

Coming now back to the interpretation of the solutions, we note that excluding the choice of $n_0 < 0$ and $n_1 > 0$ (which would correspond to a space-like naked singularity), this solution represents a naked singularity for $n_0 > 0$, $n_1 < 0$, while a horizon exists at $x_h = \left(\frac{n_1}{n_0}\right)^{n/(2n+2)}$ for a) $n_0 > 0$, $n_1 > 0$ and b) $n_0 < 0$, $n_1 < 0$. For case a), n_0 is always positive for $\Lambda < 0$, but also for $0 < \Lambda < 2\alpha^2 e^{2n\zeta_0}$. For b) we have to choose Λ positive with $\Lambda > 2\alpha^2 e^{2n\zeta_0}$. Moreover, the seize associated with the extra dimension- manifold diverges at infinity. A similar behaviour of the solutions was observed in a model without gauge fields [16]. The solutions thus correspond to black holes (if n_0 and n_1 are chosen appropriately) with toroidal horizon (note the constant seize of the 2-spheres) which is extended non-trivially into n extra dimensions.

The temperature T of static black hole solutions is given by:

$$\tilde{T} = \frac{1}{2\pi}\sqrt{-\frac{1}{4}g^{tt}g^{MN}\left(\partial_M g_{tt}\right)\left(\partial_N g_{tt}\right)} \ . \quad (29)$$

Evaluation at the horizon x_h then gives for the black hole solutions studied here (with $T = \frac{\tilde{T}}{ev}$):

$$T = \frac{a_0 n_1(n+1)}{2\pi n}x_h^{\frac{n}{2}-\frac{2}{n}-2} \ . \quad (30)$$

As can be easily checked, the derivative of T with respect to x_h, $\frac{\partial T}{\partial x_h}$, changes sign for $n = -1$ and $n = 2 \pm 2\sqrt{2}$. Since we are only interested in positive n here, we find that $\frac{\partial T}{\partial x_h} < 0$ for $n \leq 4$ and $\frac{\partial T}{\partial x_h} > 0$ for $n \geq 5$.

For the choice of the parameters $a_0 = 1$, $\zeta_0 = 0$ and $n_1 = 1$, we plot $2\pi T$ (which is actually equal to the surface gravity) over x_h for different values of n in Fig. 1.

III. AN EFFECTIVE 4-DIMENSIONAL MODEL

As in the 5-dimensional case [9] (in our notation $n = 1$) the equations given in the previous section can equally well be derived from an effective 4-dimensional Einstein-Yang-Mills-Higgs-dilaton (EYMHD) Lagrangian. In this section, we will discuss the following matter Lagrangian:

$$\begin{aligned} L_M = & -\frac{1}{4}e^{2\omega\Gamma}F^a_{\mu\nu}F^{a,\mu\nu} - \sum_{k=1}^{n}\frac{1}{2}e^{-4\omega\Psi_k}D_\mu\Phi^a_k D^\mu\Phi^a_k \\ & -\frac{1}{2}\left(\sum_{k=1}^{n}\partial_\mu\Psi_k\partial^\mu\Psi_k + \frac{2}{3}\sum_{k>k'}\partial_\mu\Psi_k\partial^\mu\Psi_{k'}\right) - \frac{\tilde{\Lambda}}{2}e^{-2\omega\Gamma} \end{aligned} \tag{31}$$

with

$$\Gamma = \sum_{k=1}^{n}\Psi_k \ . \tag{32}$$

This matter Lagrangian provides a natural extension of the $n = 1$ effective matter Lagrangian.

The kinetic part in the dilaton fields could be diagonalized, however, we find it more convenient to leave it in the form above which reveals the symmetry $\Psi_k, \Phi_k \leftrightarrow \Psi_{k'}, \Phi_{k'}$. The n-dimensional cosmological constant leads, through dimensional reduction, to a Liouville-type potential in the dilaton with a coupling constant $\tilde{\Lambda}$ related to the n-dimensional coupling constant by means of $\Lambda_{(4+n)} = 2\alpha^2\tilde{\Lambda}$. The dilatons in this theory thus become massive.

The Lagrangian (31) is then coupled minimally to Einstein gravity according to the full action

$$\begin{aligned} S &= S_G + S_M \\ &= \int\sqrt{-g^{(4)}}\left(L_G + L_M\right)d^4x \end{aligned} \tag{33}$$

where $L_G = R/(16\pi G_4)$, R is the Ricci scalar and G_4 is the 4-dimensional Newton's constant.

Note that the dilaton fields are coupled by an independent coupling constant ω to the gauge and Higgs fields. In this respect, the dilatons here are treated as independent scalar fields, while in the action (1) they appear as parts of the metric tensor.

After the rescaling

$$\Psi_k = v\psi_k \quad , \quad \omega = \frac{\gamma}{v} \quad , \quad \alpha = v\sqrt{G_4} \tag{34}$$

the resulting set of equations only depends on the coupling constants α and γ. We refrain from giving the explicit form of the equations here, but refer the reader to [11] for the case $n = 1$.

Note that the equations (14), (15), (17), (18), (19) become equivalent to the field equations associated to (33) by using the same Ansätze for the 4-dimensional metric, the gauge and Higgs fields, but by identifying

$$\zeta_k = 2\gamma\psi_k = 2\omega\Psi_k \quad , \quad \alpha^2 = 3\gamma^2 \ . \tag{35}$$

Remarkably, this identification turns out to be independent on n.

Note also that the two models differ for $n > 1$ with respect to the constraint (20) that doesn't exist in the 4-dimensional effective model.

A. Non-abelian solutions with diverging Higgs fields

So-called $AdS_{n+2} \times S^2$ solutions were first discussed for $n = 1$ and $\Lambda = 0$ in [9]. They can be thought of as hypertubes with spherically symmetric cross-section. The generalisation of these solutions is not possible for $n > 1$, since the Higgs fields don't fulfill the constraints (20).

However, we can construct the counterparts of these solutions in the 4-dimensional effective action, since the constraints don't appear here.

First, we assume that $H_1(x) = H_2(x) = ... \equiv H(x)$ and thus $\zeta_1(x) = \zeta_2(x) = ... \equiv \zeta(x)$. Further, we choose $\alpha^2 = 3\gamma^2$ to "mimick" the $4+n$-dimensional case as best as possible.

The fields have the form

$$N(x) = N_0 \quad , \quad A(x) = x^a \quad , \quad \psi_1(x) = \psi_2(x) = \ldots = \psi(x) = \frac{\sqrt{3}}{n\alpha} \ln\left(\frac{x}{R}\right) \tag{36}$$

$$K = \sqrt{q} \quad , \quad H_1(x) = H_2(x) = \ldots = H(x) = h\left(\frac{x}{R}\right)^{2/n} \tag{37}$$

The 00-component of the energy-momentum tensor for this soltuion reads:

$$T_0^0 = -\left(\frac{2N_0 h^2}{n} + nqh^2 + \frac{(q-1)^2}{2R^4}\right) \ . \tag{38}$$

This is constant, however, since $\sqrt{-g} = R^2 x^a \sin\theta$, the energy of this solution is not finite. Furthermore, the definition of a magnetic charge cannot be done in the usual sense since the Higgs field doesn't tend to its vacuum expectation value asymptotically.

Substituting this ansatz into the equations leads to a set of five algebraic equations which allow, in principle, to determine the parameters N_0, a, R, h, q. The following relations are easy to obtain:

$$a = \frac{2+n+2\kappa h^2}{n} \quad , \quad h^2 = \frac{1-q}{nR^2} \quad , \quad N_0 = \frac{qR^2n^3}{2n^2R^2 + 2\kappa(1-q)} \quad \text{with} \quad \kappa \equiv 2\alpha^2 \tag{39}$$

and leave us with a system of equations for R and q. It turns out that the solution is relatively simple in the case $n = 2$, however much more involved for $n \neq 2$. This is surely related to the fact that the Higgs field increases linearly for $n = 2$.

For $n = 2$ we find

$$R^2 = \kappa \frac{4(1-q)(1-9q^2) - \tilde{\Lambda}(1-q)^3}{48q(1+3q) + \tilde{\Lambda}(1-q)(3-8q)} \quad \text{with} \quad \tilde{\Lambda} = \kappa\Lambda \ , \tag{40}$$

and the relation determining q is given by the polynomial equation

$$\tilde{\Lambda}^2(1-q)^4 + \tilde{\Lambda}(1-q)(17-47q+4q^2-24q^3) + 16(1+3q)(1-12q+6q^2-9q^3) = 0 \ . \tag{41}$$

We solved this equation for q numerically by choosing several values of $\tilde{\Lambda}$ in $[-1, +1]$. It turns out that for the values of $\tilde{\Lambda}$ considered, only one positive solution of q exists. Then the various parameters characterizing the solutions can be computed. They are presented in Fig.2 as functions of $\tilde{\Lambda}$. As can be seen, both q and N_0 tend to zero for $\Lambda \rightarrow \frac{1}{2\alpha^2}$. This tells us that the solutions don't exist for arbitrary values of the cosmological constant. Specifically, they exist for all values of the cosmological constant as long as it is chosen to be negative (which corresponds to $(4+n)$-dimensional Anti-de Sitter space), but for positive cosmological constant only under the restriction that $\Lambda_{(4+n)} < \frac{e^2}{2G_{(4+n)}}$.

For generic values of n, the solution is more envolved. We find :

$$R^2 = \kappa \frac{P}{Q} \tag{42}$$

with

$$P = n^2(q-1)(2n^2q-2nq^2+5nq-n-2n^2+6n)(2nq+n+2q)+\tilde{\Lambda}(n^3-2n^2q-2nq+2q)(2q-n)(q-1)^2 \tag{43}$$

and

$$\begin{aligned} Q &= n^2(2n^4q - 2n^3q^2 + 5n^3q - n^3 - 6n^2q^2 + 6n^2q - 4nq^2 + 2nq - 4q)(2nq + n + 2q) \\ &+ \tilde{\Lambda}n(q-1)(-n^5 + 4n^4q - 4n^3q^2 + 4n^3q - 8n^2q^2 - 4n^2q - 2nq + 8q^2 + 2q) \ . \end{aligned} \tag{44}$$

The equation for q is a lengthly polynomial expression of degree nine in q and of degree three in $\tilde{\Lambda}$, that's why we don't give it here.

The solution presented above is completely symmetric under the exchange

$$H_j, \zeta_j \leftrightarrow H_k, \zeta_k \quad , \quad \forall j, k \in \{1, \ldots, n\} \ . \tag{45}$$

It can be generalized in the following way :

$$N(x) = N_0 \quad , \quad A(x) = x^a \quad , \quad K(x) = \sqrt{q} \tag{46}$$

and no-equal dilaton and Higgs fields

$$\psi_k(x) = \frac{\sqrt{3}}{n\alpha} \ln\left(\frac{x}{R_k}\right) \quad , \quad H_k(x) = h\left(\frac{x}{R_k}\right)^{2/n} \quad , \quad k = 1, ..., n \quad , \tag{47}$$

where $R_1, \ldots, R_n$ are positive constants which fulfill the constraint $\prod_{k=1}^{n} R_k = R^n$, where R is the quantity determined in the symmetric case. This choice of functions leads to a family of $n-1$ possible solutions.

IV. CONCLUSIONS

Adopting the dimensional reduction scheme based on the fact that the fields do not depend on the n extra dimensions, we have studied the classical equations corresponding to the SU(2) Einstein-Yang-Mills model in $4+n$ dimensions including a cosmological constant.

Supplementing the initial model with a cosmological constant results in a self-interacting potential of the dilatons, where this potential is completely fixed in terms of the coupling constants of the model.

A spherically symmetric Ansatz for the fields leads to a system of non-linear differential equations. Constraints on the fields lead to the fact that only solutions which become effectively 5-dimensional (with one non-zero Higgs fields) or with all Higgs fields constant and zero gauge fields exist. We construct the “embedded” abelian solutions with all Higgs fields constant. The Ricci and Kretschman scalars of these solutions are finite everywhere apart from the origin. Depending on the choice of parameters they represent either naked singularities or black holes. They generalize solutions obtained in [13] to the case of a $(4+n)$-dimensional space-time with an $SO(3)$ symmetry. These solutions are not asymtotically flat, neither de Sitter nor Anti-de Sitter, but all components of the energy-momentum tensor remain finite asymptotically.

One interesting generalisation of this result would be to construct the rotating counterparts of these solutions, following [17] for the Einstein-Maxwell case.

The second family of solutions obtained in this paper is the family of solutions of the effective 4-dimensional action. These solutions are characterized by non-trivial matter fields, especially diverging Higgs fields. They depend smoothly on the cosmological constant. We find that for $n = 2$ these type of solutions exist only up to a maximal value of the cosmological constant.

Acknowledgements Y. B. gratefully acknowledges the Belgian F.N.R.S. for financial support.

---

[1] T. Kaluza, Sitzungsber. Preuss. Akad. Wiss. Berlin (1921).

[2] O. Klein, Z. Phys. **37** (1926), 895.

[3] G. Gibbons and K. Maeda, Nucl. Phys. **B298** (1988), 741; D. Garfinkle, G. Horowitz and A. Strominger, Phys. Rev. **D43** (1991), 371.

[4] see e.g. J. Polchinski, *String Theory*, Cambridge University press (1998).

[5] K. Akama, “Pregeometry”, in Lecture Notes in Physics, 176, Gauge Theory and Gravitation, Proceedings, Nara, 1982, edited by K. Kikkawa, N. Nakanishi and H. Nariai, 267-271 (Springer-Verlag, 1983) [hep-th/0001113]; V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. **125 B**

(1983), 136; **125 B** (1983), 139; G. Davli and M. Shifman, Phys. Lett. **B 396** (1997), 64; **407** (1997), 452; I. Antoniadis, Phys. Lett. **B 246** (1990), 377; N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B 429** (1998), 263; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B 436** (1998), 257; L. Randall and R. Sundrum, Phys. Rev. Lett. **83** (1999), 3370; **83** (1999), 4690.

[6] J. Maldacena, Adv. Theor. Phys. **2** (1998), 231; E. Witten, Adv. Theor. Math. Phys. **2** (1998), 253; *for a review see e.g.* O. Aharony, S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, Phys. Rep. **323** (2000), 183.

[7] A. Strominger, JHEP **10** (2001), 034; JHEP **11** (2001), 049.

[8] Y. Brihaye, A. Chakrabarti, B. Hartmann and D. H. Tchrakian, Phys. Lett. **B 561** (2003), 161.

[9] M. S. Volkov, Phys. Lett. **B524** (2002), 369.

[10] Y. Brihaye, F. Clement and B. Hartmann, *Spherically symmetric Yang-Mills solutions in a* $(4+n)$*-dimensional space-time*, hep-th/0403041.

[11] B. Hartmann, Y. Brihaye and B. Bertrand, Phys. Lett. **B 570** (2003), 137.

[12] S. Poletti and D. Wiltshire, Phys. Rev. **D 50** (1994), 7260.

[13] K. C. K. Chan, J. H. Horne and R. B. Mann, Nucl. Phys. **B447** (1995), 441.

[14] K. Lee, V.P. Nair and E.J. Weinberg, Phys. Rev. **D45** (1992), 2751;
P. Breitenlohner, P. Forgacs and D. Maison, Nucl. Phys. **B383** (1992), 357;
P. Breitenlohner, P. Forgacs and D. Maison, Nucl. Phys. **B442** (1995), 126.

[15] G. ‘t Hooft, Nucl. Phys. **B79** (1974), 276;
A. M. Polyakov, JETP Lett. **20** (1974), 194.

[16] D. Wiltshire, Phys. Rev. **D 44** (1991), 1100.

[17] T. Ghosh and P. Mitra, Class. Quantum Grav. **20** (2003), 1403.



FIG. 1: The surface gravity of the embedded abelian black hole solutions, which is equal to $2\pi T$, is shown in dependence on the horizon value x_h for $n = 1, ..., 7$.



FIG. 2: The values of the constants of the non-abelian solutions of the 4-dimensional effective model (resulting for the 4 + 2-dimensional model) are shown in dependence on the cosmological constant.